

# MP469: Laplace's Equation in Spherical Polar Co-ordinates

For many problems involving Laplace's equation in 3-dimensions

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} + \frac{\partial^2 u}{\partial z^2} = 0. \quad (1)$$

it is more convenient to use spherical polar co-ordinates  $(r, \theta, \phi)$  rather than Cartesian co-ordinates  $(x, y, z)$ . These are related to each other in the usual way by

$$\begin{aligned} x &= r \cos \phi \sin \theta \\ y &= r \sin \phi \sin \theta \\ z &= r \cos \theta. \end{aligned}$$

To translate (1) into a differential equation involving  $(r, \theta, \phi)$  we need the following partial derivatives:

$$\begin{aligned} \frac{\partial x}{\partial r} &= \cos \phi \sin \theta, & \frac{\partial y}{\partial r} &= \sin \phi \sin \theta, & \frac{\partial z}{\partial r} &= \cos \theta, \\ \frac{\partial x}{\partial \theta} &= r \cos \phi \cos \theta, & \frac{\partial y}{\partial \theta} &= r \sin \phi \cos \theta, & \frac{\partial z}{\partial \theta} &= -r \sin \theta, \\ \frac{\partial x}{\partial \phi} &= -r \sin \phi \sin \theta, & \frac{\partial y}{\partial \phi} &= r \cos \phi \sin \theta, & \frac{\partial z}{\partial \phi} &= 0. \end{aligned}$$

Using these the chain rule for differentiation implies that

$$\begin{aligned} \frac{\partial}{\partial r} &= \frac{\partial x}{\partial r} \frac{\partial}{\partial x} + \frac{\partial y}{\partial r} \frac{\partial}{\partial y} + \frac{\partial z}{\partial r} \frac{\partial}{\partial z} = \cos \phi \sin \theta \frac{\partial}{\partial x} + \sin \phi \sin \theta \frac{\partial}{\partial y} + \cos \theta \frac{\partial}{\partial z}, \\ \frac{\partial}{\partial \theta} &= \frac{\partial x}{\partial \theta} \frac{\partial}{\partial x} + \frac{\partial y}{\partial \theta} \frac{\partial}{\partial y} + \frac{\partial z}{\partial \theta} \frac{\partial}{\partial z} = r \cos \phi \cos \theta \frac{\partial}{\partial x} + r \sin \phi \cos \theta \frac{\partial}{\partial y} - r \sin \theta \frac{\partial}{\partial z}, \\ \frac{\partial}{\partial \phi} &= \frac{\partial x}{\partial \phi} \frac{\partial}{\partial x} + \frac{\partial y}{\partial \phi} \frac{\partial}{\partial y} + \frac{\partial z}{\partial \phi} \frac{\partial}{\partial z} = -r \sin \phi \sin \theta \frac{\partial}{\partial x} + r \cos \phi \sin \theta \frac{\partial}{\partial y}. \end{aligned}$$

These can be inverted, by taking linear combination with trigonometric function for example, to express partial derivatives of Cartesian co-ordinates in terms of polar co-ordinates:

$$\begin{aligned} \frac{\partial}{\partial x} &= \cos \phi \sin \theta \frac{\partial}{\partial r} + \frac{\cos \phi \cos \theta}{r} \frac{\partial}{\partial \theta} - \frac{\sin \phi}{r \sin \theta} \frac{\partial}{\partial \phi} \\ \frac{\partial}{\partial y} &= \sin \phi \sin \theta \frac{\partial}{\partial r} + \frac{\sin \phi \cos \theta}{r} \frac{\partial}{\partial \theta} + \frac{\cos \phi}{r \sin \theta} \frac{\partial}{\partial \phi} \\ \frac{\partial}{\partial z} &= \cos \theta \frac{\partial}{\partial r} - \frac{\sin \theta}{r} \frac{\partial}{\partial \theta}. \end{aligned} \quad (2)$$

One way of deriving the Laplacian in 3-dimensional spherical polars is to expand the unit vectors  $\hat{\mathbf{r}}$ ,  $\hat{\boldsymbol{\theta}}$  and  $\hat{\boldsymbol{\phi}}$  in the directions of increasing  $r$ ,  $\theta$  and  $\phi$  respectively, in terms of the Cartesian unit vectors  $\hat{\mathbf{x}}$ ,  $\hat{\mathbf{y}}$  and  $\hat{\mathbf{z}}$ :

$$\begin{aligned} \hat{\mathbf{r}} &= \sin \theta \cos \phi \hat{\mathbf{x}} + \sin \theta \sin \phi \hat{\mathbf{y}} + \cos \theta \hat{\mathbf{z}}, \\ \hat{\boldsymbol{\theta}} &= \cos \theta \cos \phi \hat{\mathbf{x}} + \cos \theta \sin \phi \hat{\mathbf{y}} - \sin \theta \hat{\mathbf{z}}, \\ \hat{\boldsymbol{\phi}} &= -\sin \phi \hat{\mathbf{x}} + \cos \phi \hat{\mathbf{y}}. \end{aligned} \quad (3)$$

These can be inverted to give

$$\begin{aligned}\hat{\mathbf{x}} &= \sin \theta \cos \phi \hat{\mathbf{r}} + \cos \theta \cos \phi \hat{\boldsymbol{\theta}} - \sin \phi \hat{\boldsymbol{\phi}}, \\ \hat{\mathbf{y}} &= \sin \theta \sin \phi \hat{\mathbf{r}} + \cos \theta \sin \phi \hat{\boldsymbol{\theta}} + \cos \phi \hat{\boldsymbol{\phi}}, \\ \hat{\mathbf{z}} &= \cos \theta \hat{\mathbf{r}} - \sin \theta \hat{\boldsymbol{\theta}}.\end{aligned}\tag{4}$$

Then define the vector differential operator

$$\nabla := \hat{\mathbf{x}} \frac{\partial}{\partial x} + \hat{\mathbf{y}} \frac{\partial}{\partial y} + \hat{\mathbf{z}} \frac{\partial}{\partial z}\tag{5}$$

and the Laplacian can be written as  $\nabla^2 = \nabla \cdot \nabla$ , using the vector dot product. Putting (2) and (4) in (5), and using standard trigonometric identities, gives

$$\nabla := \hat{\mathbf{r}} \frac{\partial}{\partial r} + \hat{\boldsymbol{\theta}} \frac{1}{r} \frac{\partial}{\partial \theta} + \hat{\boldsymbol{\phi}} \frac{1}{r \sin \theta} \frac{\partial}{\partial \phi}.\tag{6}$$

Now to calculate  $\nabla^2 = \nabla \cdot \nabla$  in spherical polars we must be careful since the polar unit vectors  $\hat{\mathbf{r}}$ ,  $\hat{\boldsymbol{\theta}}$  and  $\hat{\boldsymbol{\phi}}$  are not constant. From (3)

$$\begin{aligned}\frac{\partial}{\partial r} \hat{\mathbf{r}} &= 0, & \frac{\partial}{\partial r} \hat{\boldsymbol{\theta}} &= 0, & \frac{\partial}{\partial r} \hat{\boldsymbol{\phi}} &= 0, \\ \frac{\partial}{\partial \theta} \hat{\mathbf{r}} &= \hat{\boldsymbol{\theta}}, & \frac{\partial}{\partial \theta} \hat{\boldsymbol{\theta}} &= -\hat{\mathbf{r}}, & \frac{\partial}{\partial \theta} \hat{\boldsymbol{\phi}} &= 0, \\ \frac{\partial}{\partial \phi} \hat{\mathbf{r}} &= -\sin \theta \hat{\boldsymbol{\phi}}, & \frac{\partial}{\partial \phi} \hat{\boldsymbol{\theta}} &= \cos \theta \hat{\boldsymbol{\phi}}, & \frac{\partial}{\partial \phi} \hat{\boldsymbol{\phi}} &= -\sin \theta \hat{\mathbf{r}} - \cos \theta \hat{\boldsymbol{\theta}}.\end{aligned}$$

Using these in (6) the Laplace differential operator in equation (1) can be expressed directly in terms of spherical polar co-ordinates:

$$\begin{aligned}\nabla^2 u &= \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} + \frac{\partial^2 u}{\partial z^2} \\ &= \frac{1}{r} \frac{\partial^2 (ru)}{\partial r^2} + \frac{1}{r^2 \sin \theta} \frac{\partial}{\partial \theta} \left( \sin \theta \frac{\partial u}{\partial \theta} \right) + \frac{1}{r^2 \sin^2 \theta} \frac{\partial^2 u}{\partial \phi^2}.\end{aligned}$$