

# Mathematical Physics MP202: Mechanics

## Problem Sheet 5

24th April. 2007

1) A system of  $N$  particles with masses  $m_i$  at points  $\mathbf{r}_i$ ,  $i = 1, \dots, N$ , and total mass  $M = \sum_{i=1}^N m_i$  consists of two disjoint parts with masses  $M_1$  and  $M_2$  ( $M = M_1 + M_2$ ) and centres of mass  $\mathbf{R}_1$  and  $\mathbf{R}_2$  respectively. Show that the centre of mass of the whole system is

$$\mathbf{R} = \frac{M_1 \mathbf{R}_1 + M_2 \mathbf{R}_2}{M}.$$

Find the centre of mass of the Earth-Moon system, relative to the centre of the Earth, given that mass of the Earth is  $M_1 = 6.0 \times 10^{24} \text{ kg}$ , the mass of the Moon is  $M_2 = 7.3 \times 10^{22} \text{ kg}$  and the distance between the Earth and the Moon is  $3.8 \times 10^8 \text{ m}$ . How does your answer compare with the radius of the Earth, which is  $6,400 \text{ km}$ ?

2) Calculate the moment of inertia of a flat circular disc of mass  $M$  and radius  $a$  about the centre of the disc. The disc is pierced half-way between its centre and the edge and suspended in a vertical plane from a frictionless pivot. Calculate the period of small oscillations of the disc about its equilibrium position in terms of  $M$ ,  $a$  and the acceleration due to gravity,  $g$ .

3) Show that the kinetic energy of a particle of mass  $m$  moving in a two-dimensional plane, when expressed in two-dimensional polar co-ordinates, is

$$\frac{1}{2}m(\dot{r}^2 + r^2\dot{\phi}^2).$$

Using  $r$  and  $\phi$  as generalised co-ordinates determine the generalised momenta associated with them and derive the Euler-Lagrange equations of motion for  $m$ . Solve the equations for an arbitrary initial position and velocity and show that the motion is a straight line.

4) A bead of mass  $m$  is constrained to move on a frictionless wire with shape  $y(x) = a \cos(bx)$  in a vertical plane, with a constant gravitational field and  $y$  measured vertically upwards ( $a$  and  $b$  are positive constants). Find the Lagrangian for the bead and derive the equations of motion.

5) Two masses  $m_1$  and  $m_2$  are hung around a frictionless pulley by a light inelastic cord of length  $l$ , each hanging vertically downwards. Ignoring any sideways motion of the mass so they only move vertically and ignoring the mass of the pulley, find the Lagrangian for the system and derive the equations of motion.

6) A bead of mass  $m$  slides on a frictionless wire in the shape of a circle of radius  $a$ , rotating with constant angular velocity  $\omega$  about an axis in the same plane as the wire and passing through its centre. The position of the bead is given by

$$x = \sin \psi \cos \omega t, \quad y = \sin \psi \sin \omega t, \quad z = a(1 - \cos \psi),$$

where  $0 \leq \psi \leq \pi$ .

Using  $\psi$  as a generalised co-ordinate show that the Lagrangian is

$$L = \frac{m}{2}a^2\dot{\psi}^2 - V_{eff}(\psi)$$

where the effective potential is

$$V_{eff}(\psi) = -mga \cos \psi - \frac{1}{2}ma^2\omega^2 \sin^2 \psi.$$

Sketch the effective potential for  $0 \leq \omega^2 < g/a$  and  $\omega^2 > g/a$  and identify points of stable and unstable equilibrium in both cases.