

# MP 472 Quantum Information and Computation

<http://www.thphys.may.ie/staff/jvala/MP472.htm>

## Outline

Open quantum systems

The density operator

Quantum noise (decoherence)

- Quantum operations
- Operator sum representation

Quantum error correction

Fault-tolerant quantum  
computation

## Operator sum representation (review)

is a representation of quantum operations in terms of the operators on the (principal) system only (important!):

Let  $|e_k\rangle$  be the orthonormal basis for the (finite dimensional) Hilbert space of the environment, and let  $\rho = |e_0\rangle\langle e_0|$  be the initial (pure) state of the environment. Then we can express quantum operations as

$$\mathcal{E}(\rho) = \sum_k \langle e_k|U(\rho \otimes |e_0\rangle\langle e_0|)U^\dagger|e_k\rangle = \sum_k E_k \rho E_k^\dagger$$

where  $E_k = \langle e_k|U|e_0\rangle$  is an operator on the Hilbert space of the (principal) system; It is called an operation element of quantum operation.

The operation elements satisfy an important constraint known as **the completeness relation**

$$\sum_k E_k^\dagger E_k = I$$

The completeness relation is satisfied by quantum operations  $\mathcal{E}(\rho)$  which are trace-preserving.

In general, there are non-trace-preserving operations for which  $\sum_k E_k^\dagger E_k \leq I$  but they describe processes in which extra information about what occurred in the process is obtained by measurement.

## Physical interpretation of OSR (review)

Imagine that a measurement of the environment is performed in the basis  $|e_k\rangle$  after the unitary operation  $U$  has been applied. By the principle of implicate measurement (Notes 7), such a measurement affects only the state of the environment.

Let  $\rho_k$  be the state of the principal system given that outcome  $k$  occurs, so

$$\rho_k \propto \text{tr}_E(|e_k\rangle\langle e_k|U(\rho \otimes |e_0\rangle\langle e_0|)U^+|e_k\rangle\langle e_k|) = \langle e_k|U(\rho \otimes |e_0\rangle\langle e_0|)U^+|e_k\rangle = E_k\rho E_k^+$$

Normalizing  $\rho_k$

$$\rho_k = E_k\rho E_k^+ / \text{tr}(E_k\rho E_k^+)$$

The probability of outcome  $k$  is given by

$$p(k) = \text{tr}(|e_k\rangle\langle e_k|U(\rho \otimes |e_0\rangle\langle e_0|)U^+|e_k\rangle\langle e_k|) = \text{tr}(E_k\rho E_k^+)$$

Thus

$$\mathcal{E}(\rho) = \sum_k p(k) \rho_k = \sum_k E_k\rho E_k^+$$


The action of the quantum operation is equivalent to taking the state  $\rho$  and randomly replacing it by

$$\rho_k = E_k\rho E_k^+ / \text{tr}(E_k\rho E_k^+)$$

with probability

$$p_k = \text{tr}(E_k\rho E_k^+)$$

## Physical interpretation of OSR: example (review)



Suppose the states  $|e_k\rangle$  are chosen as  $|0_E\rangle$  and  $|1_E\rangle$ .

Measurement in the computational basis of the environment qubit does not change the state of the principal system:

$$U_{\text{CNOT}} = |0_S 0_E\rangle\langle 0_S 0_E| + |0_S 1_E\rangle\langle 0_S 1_E| + |1_S 1_E\rangle\langle 1_S 0_E| + |1_S 0_E\rangle\langle 1_S 1_E|$$

Thus

$$E_0 = \langle 0_E | U_{\text{CNOT}} | 0_E \rangle = |0_S\rangle\langle 0_S|$$

$$E_1 = \langle 1_E | U_{\text{CNOT}} | 1_E \rangle = |1_S\rangle\langle 1_S|$$

and therefore

$$\mathcal{E}(\rho) = P_0 \rho P_0 + P_1 \rho P_1$$

This result is in agreement with the result of our previous example (see previous notes - slide 5).

# Measurements and OSR

How do we determine OSR for a given open quantum system?

A) Unitary dynamics

$$E_k = \langle e_k | U | e_0 \rangle$$

B) Measurement on the combined system-environment (S-E) (here  $\sum_k E_k + E_k \leq I$ )

Let an initial (product) state  $\rho = \rho_s \otimes \rho_{env}$  to evolve under unitary dynamics  $U$  and then allow (projective) measurement on S-E; the final quantum state of S-E is

$$P_m U (\rho_s \otimes \rho_{env}) U^\dagger P_m / \text{tr}(P_m U (\rho_s \otimes \rho_{env}) U^\dagger P_m)$$

The final state of S only is obtained by tracing out E:

$$\text{tr}_E (P_m U (\rho_s \otimes \rho_{env}) U^\dagger P_m) / \text{tr}(P_m U (\rho_s \otimes \rho_{env}) U^\dagger P_m)$$

Define a map

$$\mathcal{E}_m(\rho) = \text{tr}_E (P_m U (\rho_s \otimes \rho_{env}) U^\dagger P_m) = \rho_{env} = \sum_j q_j |j\rangle\langle j|$$

and using orthonormal basis  $|e_k\rangle$  for the environment we obtain

$$= \sum_{jk} q_j \langle e_k | P_m U (\rho_s \otimes \rho_{env}) |j\rangle\langle j| U^\dagger P_m |e_k\rangle = \sum_{jk} E_{jk} \rho_s E_{jk}^\dagger$$

$$E_{jk} = (q_j)^{1/2} \langle e_k | P_m U |j\rangle$$

## System-environment models for any OSR

Given  $\{E_k\}$ , is there a reasonable model environment and dynamics to produce a quantum operation with the operational elements  $\{E_k\}$ ?

For any quantum operation (trace-preserving or non-trace-preserving)  $\mathcal{E}$ , with operational elements  $\{E_k\}$ , there exist a model environment  $E$ , starting in a pure state  $|e_0\rangle$ , and model dynamics specified by a unitary operator  $U$  and projector  $P$  onto  $E$  s.t.

$$\mathcal{E}(\rho) = \text{tr}_E (PU (\rho_s \otimes |e_0\rangle\langle e_0|)U^\dagger P)$$

---

To show this, let's assume that  $\mathcal{E}$  is a trace-preserving quantum operation, with operator sum representation generated by operation elements  $\{E_k\}$ , satisfying the completeness relation. In this case, we thus need to find only an appropriate unitary operator  $U$  to model the dynamics. Let  $|e_k\rangle$  be an orthonormal basis set for environment (with one-to-one correspondence with the index  $k$  for  $E_k$ ).

Define an operator  $U$  s.t.

$$U|\psi\rangle|e_0\rangle = \sum_k E_k|\psi\rangle|e_0\rangle$$

where  $|e_0\rangle$  is some standard state of the environment. For arbitrary states of the principal system  $|\psi\rangle$  and  $|\phi\rangle$ ,  $\langle\psi|\langle e_0|U^\dagger U|\phi\rangle|e_0\rangle = \langle\psi|\phi\rangle$  (by the completeness relation) so the operator  $U$  acts unitarily on the S-E state space, and tracing the state over  $E$

$$\text{tr}_E (U (\rho_s \otimes |e_0\rangle\langle e_0|)U^\dagger) = \sum_k E_k \rho E_k^\dagger$$

shows that this model provides a realization of the quantum operation  $\mathcal{E}$  with  $\{E_k\}$ .

# Axiomatic approach to quantum operations

Abstract but powerful!

Quantum operation  $\mathcal{E}$  is defined as a map from the set of density operators of the input space  $S_1$  to the output space  $S_2$ , with the following axioms:

(A 1)

$\text{tr}[\mathcal{E}(\rho)]$  is the probability that the process represented by  $\mathcal{E}$  occurs, when  $\rho$  is the initial state; thus,  $0 \leq \text{tr}[\mathcal{E}(\rho)] \leq 1$  for any state  $\rho$ .

(A 2)

$\mathcal{E}$  is a convex-linear map on the set of density matrices, i.e. for probabilities  $\{p_i\}$ ,

$$\mathcal{E}(\sum_i p_i \rho_i) = \sum_i p_i \mathcal{E}(\rho_i).$$

(A 3)

$\mathcal{E}$  is a completely positive map. That is if  $\mathcal{E}$  maps density operators of system  $S_1$  to density operators of system  $S_2$ , then  $\mathcal{E}(A)$  must be positive for any positive operator  $A$ . Furthermore, if we introduce an extra system  $R$  of arbitrary dimensionality, it must be true that  $(\mathcal{I} \otimes \mathcal{E})(A)$  is positive for any positive operator  $A$  on the combined system  $RS_1$ , where  $\mathcal{I}$  denotes the identity map on system  $R$ .


Theorem: The map  $\mathcal{E}$  satisfies axioms A1, A2 and A3 iff

$$\mathcal{E}(\rho) = \sum_k E_k \rho E_k^\dagger$$

for some set of operators  $\{E_k\}$  which map the input Hilbert space to the output Hilbert space, and  $\sum_k E_k^\dagger E_k \leq I$ .

## Unitary freedom in OSR

The operational elements in an operator sum representation for a quantum operation are not unique.



### Theorem:

Suppose  $\{E_i, \dots, E_m\}$  and  $\{F_i, \dots, F_n\}$  are operation elements giving rise to quantum operations  $\mathcal{E}$  and  $\mathcal{F}$ , respectively. By appending zero operators to the shorter list of operational elements we may ensure that  $m=n$ . Then  $\mathcal{E}=\mathcal{F}$  iff there exist complex numbers  $u_{ij}$  s.t.  $E_i = \sum_j u_{ji} F_j$ , and  $u$  is an  $m$ -by- $m$  matrix.